

PATIENT INFORMATION

BUSPIRONE HYDROCHLORIDE TABLETS

(byoo-SPYE-rone hye" droe klor'ide)

HOW TO USE:

BusPIRone Hydrochloride Tablets

15 mg and 30 mg

Response to buspirone varies among individuals. Your physician may find it necessary to adjust your dosage to obtain the proper response.


This tablet design makes dosage adjustments easy. Each tablet is scored and can be broken accurately to provide any of the following dosages:

If your doctor prescribed the 15 mg Tablet:

Front Side:



Other Side:



10 mg (two thirds of a tablet)



5 mg (one third of a tablet)



7.5 mg (One-half of a tablet)




If your doctor prescribed 30 mg Tablet:

Front Side:



Other Side:



20 mg (two thirds of a tablet)




10 mg (one third of a tablet)



15 mg (One-half of a tablet)



To break a tablet accurately and easily, hold the tablet between your thumbs and index fingers close to the appropriate tablet score (groove) as shown below. Then, with the tablet score facing you, apply pressure and snap the tablet segments apart (segments breaking incorrectly should not be used).



Manufactured by:

Graviti Pharmaceuticals Pvt. Ltd.
Telangana-502307, INDIA.

M.L. No.: 12/SRD/TS/2017/F/G

Manufactured for:

Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Issued: 02/2025

PILR65450-00

Dispense with Medication Guide available at:

<https://risingpharma.com/Medguides/BuspironeHydrochlorideTabletsMG.pdf>